

This is a repository copy of *Alcohol complicates multimorbidity in older adults*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/148108/>

Version: Published Version

Article:

Stewart, Duncan orcid.org/0000-0001-7355-4280 and McCambridge, Jim orcid.org/0000-0002-5461-7001 (2019) Alcohol complicates multimorbidity in older adults. *BMJ*. 365:l4304. ISSN 1756-1833

<https://doi.org/10.1136/bmj.l4304>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



EDITORIALS

Alcohol complicates multimorbidity in older adults

Attention to alcohol should be integral to routine care

Duncan Stewart *senior lecturer in addictive behaviours and public health*, Jim McCambridge *professor of addictive behaviours and public health*

Department of Health Sciences, University of York, York, UK

Lower birth rates and higher life expectancy are transforming global population age profiles, such that the prevalence of long term health conditions in the general population is rising steeply, with more people living with multimorbidity as they age.¹ Most people aged 65 or over have multiple conditions,¹ and the proportion with multimorbidity can be expected to increase further. Trends in alcohol consumption show higher levels of unhealthy drinking among older birth cohorts in high income countries, and older people in the UK drink more than those in other European countries, Australasia, and North America.² The challenges of clinical management of older adults with multimorbidity are made more difficult by alcohol consumption, even at low doses.

Health risks

Drinking alcohol increases risks to health in a largely dose-response manner.³ It is a particular concern among older people because metabolic efficiency is diminished with age. As a toxic and addictive drug, alcohol can make existing health problems worse by posing direct risks of organ damage³ as well as by compromising treatment effectiveness and the safety of prescribing. For older patients prescribed medications for multiple conditions, the potential for non-adherence, adverse events, and drug interactions requires careful management.⁴

This is made more challenging by changes to the absorption, distribution, and metabolism of alcohol in older age groups, which make interactions with medications more likely.⁵ Interactions can occur across a range of common medications for long term conditions, including cardiovascular diseases, pain, depression, diabetes, musculoskeletal conditions, infections, and cancer.⁶ Medications that interact with alcohol are widely prescribed to people who drink alcohol.⁷ Alcohol can also have a direct and adverse effect on many conditions for which medications are prescribed, even when consumption is lower than that which is defined as low risk in drinking guidelines.³

The damaging effects of alcohol consumption on the health of older people in the UK are evident in higher alcohol related hospital admissions and mortality among people aged over 65 than in younger people.⁸ The situation is deteriorating, with

substantial increases in alcohol specific deaths among 55-79 year olds since 2001.⁹ As the prevalence of multimorbidity grows, the burden of alcohol consumption can be expected to increase further, particularly in more deprived communities where multimorbidity is most prevalent¹ and alcohol's health damage greatest.¹⁰ Alcohol complicates health inequalities across the lifecourse.¹

The NHS, like other health systems, is poorly designed to meet the challenges of an ageing population.¹¹ Understanding the relationships between multiple conditions in older adults is intrinsically complex, and most treatments and clinical guidelines are based on evidence from trials for single conditions, conducted among younger and fitter populations taking fewer medications than older people with multimorbidity.⁴ The Academy of Medical Sciences has identified research into risk factors associated with common clusters of long term conditions, and how these interact, as a priority.¹¹

Clinicians must contend with major gaps in the research evidence,¹¹ and there are limitations in the clinical guidance available to practitioners. Although the guidance says that identification of hazardous drinking is important, it carries no specific recommendations on how alcohol consumption should be managed in relation to potential effects on multimorbidity.⁴

Research on the complexities of multimorbidity in older adults is urgently required as health systems struggle to cope with rising demand. For example, dedicated approaches to multimorbidity evaluated in a recent Cochrane review offer little or no improvement in clinical outcomes, health service use, and medication adherence but have some benefits for mental health.¹² Support for effective interventions for unhealthy drinking in older adults is more meagre still, with mixed results from a small number of trials of low-to-moderate quality.¹³ Worldwide, there are no trials evaluating interventions that tackle drinking in older people with multimorbidity.

Missed opportunity

Progress requires that alcohol consumption becomes integral to societal and health system responses to multimorbidity, particularly in countries like the UK where alcohol is culturally

embedded but not prominent in debates about health service reform. The NHS long term plan¹⁴ provides an example of a missed opportunity to reconsider what preventing alcohol related harm means in the context of an ageing population characterised by multimorbidity.

The plan identifies alcohol consumption and other modifiable risk behaviours as standalone risk factors, and thus as discrete targets for intervention.¹⁴ The focus on hospital admissions related to alcohol dependence ignores the complexity of health problems in older adults and the need for holistic approaches to management.¹¹ Attention to the insidious ways in which alcohol is implicated in health problems is required in a system-wide perspective on how to help older people live longer and improve their quality of life.¹⁵ Integrating attention to alcohol within routine care for long term conditions could help place prevention at the heart of health systems.¹⁴ First, however, we need new thinking about prevention.

Competing interests: We have read and understood BMJ policy on declaration of interests and have no relevant interests to declare.

Provenance and peer review: Not commissioned; externally peer reviewed.

- 1 Barnett K, Mercer SW, Norbury M, Watt G, Wyke S, Guthrie B. Epidemiology of multimorbidity and implications for health care, research, and medical education: a cross-sectional study. *Lancet* 2012;380:37-43. 10.1016/S0140-6736(12)60240-2 22579043
- 2 Gell L, Meier PS, Goyder E. Alcohol consumption among the over 50s: international comparisons. *Alcohol Alcohol* 2015;50:1-10. 10.1093/alcalc/agu082 25433252
- 3 GBD 2016 Alcohol Collaborators. Alcohol use and burden for 195 countries and territories, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 2018;392:1015-35. 10.1016/S0140-6736(18)31310-2 30146330

- 4 National Institute for Health and Care Excellence. *Multimorbidity: clinical assessment and management (NG56)*. NICE, 2016.
- 5 Meier P, Seitz HK. Age, alcohol metabolism and liver disease. *Curr Opin Clin Nutr Metab Care* 2008;11:21-6. 10.1097/MCO.0b013e3282f30564 18090653
- 6 Holton AE, Gallagher PJ, Ryan C, Fahey T, Cousins G. Consensus validation of the POSAMINO (potentially serious alcohol-medication interactions in older adults) criteria. *BMJ Open* 2017;7:e017453. 10.1136/bmjopen-2017-017453 29122794
- 7 Holton AE, Gallagher P, Fahey T, Cousins G. Concurrent use of alcohol interactive medications and alcohol in older adults: a systematic review of prevalence and associated adverse outcomes. *BMC Geriatr* 2017;17:148. 10.1186/s12877-017-0532-2 28716004
- 8 Wadd S, Papadopoulos C. Drinking behaviour and alcohol-related harm amongst older adults: analysis of existing UK datasets. *BMC Res Notes* 2014;7:741. 10.1186/1756-0500-7-741 25332101
- 9 Office for National Statistics. Alcohol-specific deaths in the UK: registered in 2017. 2018 <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/causesofdeath/bulletins/alcoholrelateddeathsintheunitedkingdom/registeredind2017#alcohol-specific-death-rates-by-age-uk>
- 10 Bellis MA, Hughes K, Nicholls J, Sheron N, Gilmore I, Jones L. The alcohol harm paradox: using a national survey to explore how alcohol may disproportionately impact health in deprived individuals. *BMC Public Health* 2016;16:111. 10.1186/s12889-016-2766-x 26888538
- 11 Academy of Medical Sciences. Multimorbidity: a priority for global health research. 2018. <https://acmedsci.ac.uk/file-download/82222577>
- 12 Smith SM, Wallace E, O'Dowd T, Fortin M. Interventions for improving outcomes in patients with multimorbidity in primary care and community settings. *Cochrane Database Syst Rev* 2016;3:CD006560. <https://www.ncbi.nlm.nih.gov/pubmed/26976529>. 10.1002/14651858.CD006560.pub3 26976529
- 13 Kelly S, Olanrewaju O, Cowan A, Brayne C, LaFortune L. Interventions to prevent and reduce excessive alcohol consumption in older people: a systematic review and meta-analysis. *Age Ageing* 2018;47:175-84. 10.1093/ageing/afx132 28985250
- 14 NHS. The NHS long term plan. 2019. <https://www.longtermplan.nhs.uk/wp-content/uploads/2019/01/nhs-long-term-plan.pdf>
- 15 McCambridge J, Saitz R. Rethinking brief interventions for alcohol in general practice. *BMJ* 2017;356:j116. 10.1136/bmj.j116 28108452

Published by the BMJ Publishing Group Limited. For permission to use (where not already granted under a licence) please go to <http://group.bmj.com/group/rights-licensing/permissions>